

(Annals of KEMC)

1. Vol. 17 No. 3 (2011)

DOI: <https://doi.org/10.21649/akemu.v17i3>

Published: 02/27/2012

Research Articles

- **Precipitating factors of Hepatic Coma**

Saira Afzal, Maaz Ahmad

213

- [FULLTEXT PDF](#)

- **Urological Injuries during Obstetrical and Gynecological Surgeries**

Rafiq Ahmed Sahito, Gulshan Ali Memon, Muhammad Saleh Khaskheli

218

- [FULLTEXT PDF](#)

- **Maternal Risk Factors Associated with Low Birth Weight: A Case Control Study**

Fariha Anjum, Tahir Javed, Muhammad Faheem Afzal, Ghazanfar Ali Sheikh

223

- [FULLTEXT PDF](#)

- **Role of Routine Preoperative Biochemistry in Anaesthesia**

Muhammad Ishaq, Babar Hussain Khan, Sarmad Masud, Rehana S Kamal, Muhammad Atif Malik

229

- [FULLTEXT PDF](#)

- **Comparison of Three Different Doses of 0.75% Hyperbaric Bupivacaine for Intraoperative Analgesia in Elective Cesarean Sections**

Muhammad Ahmad Khan, Mansab Amin Chaudhary, Tahira Yasmin, Madeeha Jabeen, Parveen Iqbal

234

◦ [FULLTEXT PDF](#)

- **Chronic Lead Acetate Nephrotoxicity: A Histological Study on Albino Rats**

Mohammad Faseeh Nisar, Iqbal Nasir, Samina Shaheen, Attiya Khalid, Nadia Tazeen

239

◦ [PDF](#)

- **Pain Modulation in Frozen Shoulder – Electrotherapy Versus Exercise Therapy**

Mirza Obaid Baig, Uzma Arif Baig, Asma Naseem

245

◦ [PDF](#)

- **Comparison of Visual Outcome after Phacoemulsification Versus Extracapsular Cataract Extraction**

Chaudhry Nasir Ahmad, Ateeq Yousif, Muhammad Zahid Siddique, Hashir Amin Malik

251

◦ [PDF](#)

- **Prevalence of Cervical Abnormalities and Co-existent Human Papilloma Virus Infection in a Mixed Bahraini Population**

Fauzia Maqsood, Wasima Arif, Saima Iqbal, Zafar Iqbal Bajwa, Muhammad Ali

256

◦ [PDF](#)

- **Study to Evaluate Hepatotoxic Characterizations Induced Through Ethanol in Albino Rats Mus Musculus**

Muhammad Habib-ur- Rehman, Farooq Ahmad, Mohammad Zubair, Tauqir Ahmad, Sharif Mughal

262

◦ [PDF](#)

- **A Clinicopathological Study of Thyroid Cancers at King Edward Medical University / Mayo Hospital, Lahore**

Shahida Niazi, Madiha Arshad, Mulazim Hussain Bukhari

268

◦ [PDF](#)

- **Delays in Management of Post Partum Psychosis in Third World Countries; A Case Report and Literature Review**

Ali Madeeh Hashmi, Asad Khizar Malik

277

◦ [PDF](#)

- **Family History of Diabetes and Parental Consanguinity: Important risk for Impaired Fasting Glucose**

Adeela Shahid, **Shahid Saeed**, Sobia Rana, Saqib Mahmood

280

◦ [PDF](#)

- **Evaluation of Ligamentotaxis in Management of Comminuted Intra-articular Fractures of Distal Radius Using Spanning Nasserâ€™Awais (NA) External Fixator**

Mohammed Iqbal, Abdul Latif Sami, M. Akram, Faisal Masood, S. M. Awais, Asif Hanif

286

◦ [PDF](#)

- **Educational Environment at University Medical and Dental College, FSD**

Aneela Umer, Sadia Khan, Musarrat-ul- Hussnaian, Sumaira Ihsan

292

◦ [PDF](#)

- **Changes in Stroke Ratio in a Secondary Referral Center of Lahore District â€™Variations in Different Seasons**

Amina Husnain, Saima Nauman, Muhammad Hussain, Khalid Umar Gill, Arif Siddiqi

299

◦ [PDF](#)

- **Maternal Mortality in Services Hospital Gyane Unit II in the Year 2010**

Alia Zaineb Asad, Shazai Ashraf, Saqib Siddiq

305

◦ [PDF](#)

- **Prevalence of Hepatitis B and C Virus Among Blood Donors in Sukkur**

Aftab Ahmed Soomro, Abdul Khaliq Abro, Bahawaldin Jamro

309

◦ [PDF](#)

- **Influence of Different Sociodemographic Factors on Body Mass Index - A Cross Sectional Survey from KEMU, Lahore**

Zahra Azeem, Yasmeen Azeem, Asif Hanif, Riaz Ahmed Warraich

314

◦ [PDF](#)

- **Complications and Management of Unsafe Abortion**

Abida Sajid, Samia Malik

320

◦ [PDF](#)

- **7 Year Follow up of Renal Artery Stenosis due to Fibromuscular Dysplasia**

Bilqees Akhtar, Muhammad Khawar Saeed, Faryal Siddique

324

◦ [PDF](#)

Editorial

- **Problems of Medical Education in Pakistan**

Syed Muhammad Awais

209

- [FULLTEXT PDF](#)
- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)
- [Open Journal Systems](#)
- [Make a Submission](#)
- [Browse](#)

ISSN : 2079-0694 (Online) | 2079-7192(Print) Previous ISSN : 1684-6680 (Annals of KEMC)

Developed By: IT Department, King Edward Medical University

© Platform & Workflow by: Open Journal Systems
Designed by Material Theme

Original Article

Family History of Diabetes and Parental Consanguinity: Important risk for Impaired Fasting Glucose

Adeela Shahid,¹ Shahid Saeed,² Sobia Rana,³ Saqib Mahmood⁴

Abstract

Objective: Offspring of type 2 diabetics have increased risk of metabolic disturbances. The aim of the

study was to access the potential effect of family history of type 2 diabetes (FHD) and parental consanguinity on fasting plasma glucose (FPG) levels.

Design: Non diabetic offspring of one or both parents

with type 2 diabetes and healthy controls of comparable age, without a FHD were the subjects of this study. FHD was defined by the presence of type 2 diabetes in one or both parents of the subject. Consanguinity was defined as history of marriage with first cousin. FPG levels were determined in cases and controls.

Shahid A.¹
Assistant Professor
Department of Physiology and Cell Biology
Shalamar Medical and Dental College, Shalamar Link Road
Lahore

Saeed S.²
Department of Physiology
King Edward Medical University, Lahore

Rana S.³
Department of Human Genetics and Molecular Biology
University of Health Sciences
Lahore

Mahmood S.⁴
Department of Human Genetics

University of Health Sciences, Lahore

Results: IFG was identified in 42% subjects with a FHD and in 14% without FHD. We found a strong independent association of FHD with impaired fasting glucose in both males and females by logistic regression analysis after adjusting the data for age sex and BMI. Parental consanguinity modifies the effect of FHD on IFG.

Conclusion: We concluded that family history of diabetes and parental history of consanguinity determine the risk for impaired fasting glucose in our population.

Keywords: Impaired fasting glucose; Parental consanguinity; Family history of diabetes.

Introduction

Type 2 diabetes mellitus (T₂DM) is the predominant form of diabetes accounting for 90% of all cases paralleling an increase in the incidence of obesity.^{1,2} It is one of the major public health challenges of 21st century and its global prevalence has attained an epidemic proportion.³ Recent increase in the prevalence of diabetes reflects the environmental, behavioral and life style changes with consumption of high caloric diet and reduced physical activity. Previously T₂DM was a disease of middle aged and older people, usually presents at age > 40 years. In the recent decades the age of onset of T₂DM has decreased and increasingly it is being observed at a younger age.⁴ Much attention has been paid in the last few years on the identification of individuals at risk for developing T₂DM.

T₂DM is a multifactorial and heterogeneous condition due to complex interaction of genetic and environmental factors. The high incidence of T₂DM among first degree relatives, a high concordance in identical twins and the increased prevalence in certain ethnic groups, provide strong evidence that genetic factors underlie the susceptibility to diabetes^{5,6}. Genetic background of diabetes increases the risk of metabolic abnormalities.^{7,8} Evidence is available suggesting disturbances of carbohydrate and lipid metabolism in individuals with a history of T₂DM in first degree relatives.⁹⁻¹² Previous studies reported higher fasting plasma glucose (FPG) in non-diabetic males with a FHD as compared to those with no FHD.^{13,14} A study reported that Asian Indian adolescents with both diabetic parents had high FPG as compared to one diabetic parent.¹⁵ While another study failed to report any difference of FPG in subjects with FHD as compared to the subjects without FHD.¹⁶ Some of the studies also reported the association between FHD and hyperglycemia.^{17,18} IFG is associated with cardiovascular risk factors, such as hypertension and dyslipidaemia¹⁹. Impaired fasting glucose (IFG) has been reported in 19.1% of Mexican children and adolescents demonstrating a high prevalence of Prediabetes in these subjects.²⁰ Moreover the presence of FHD in Mexican children and adolescents associated with IFG.²¹ Obesity is also considered a contributing factor for T₂DM. In overweight Latino children with FHD increased risk of impaired glucose tolerance test has also been observed.²²

The association of FPG with BMI have been reported in children in some studies while others do not support this association.^{23,24,13} Furthermore studies in different populations and geographical areas demonstrate an increase susceptibility of certain ethnic groups to develop T₂DM. The consanguineous marriages are quite common in Pakistan (South East Asia), with first cousin marriages being the most common. Consanguinity along with cultural and environmental factors may be responsible for early onset of the disease in our population. Keeping in view the genetic, ethnic and cultural variations, we investigated the interaction between FHD and parental consanguinity on IFG in Pakistani subjects.

Materials and Methods

Study Design

It was a cross – sectional study.

Setting

Study was carried out in the Center for Research in

Endocrinology and Reproductive Sciences (CRERS)
University of Health Sciences Lahore Pakistan.

Selection and Description of Participants

A total of 226 subjects of both sexes between 10 – 25 years of age were included in the study. Subjects were divided in the following 2 groups:

Group I: Subjects with a history of T₂DM in one or both parents (n = 124).

Group II: Subjects with no history of T₂DM in any parent (n = 102).

A total of 500 known diabetic parents from diabetic centers and clinics in Lahore were enrolled for the study. Male and female offspring (n = 124, age 10 – 25 years) of (one or both) diabetic parents volunteered and were selected to participate in the study. Non diabetic friends, neighbors and distant relatives of diabetic parents were also enrolled for the control group. Age matched 102 offspring of non diabetic parents volunteered and included in the study. Subjects of 15 – 25 years were included to examine the effect of FHD and parental consanguinity at a younger age group and more over the age for the onset of T₂DM has also decreased in the recent decade. FHD was defined by the presence of T₂DM in one or both parents of the subject.

Written informed consent from the subjects / and or parents was obtained before administering the questionnaire. Questionnaire was completed regarding the detailed FHD, medical history, past history and history of parental consanguinity. Presence of T₂DM in either parent was ascertained by detailed history, medical examination, laboratory investigations (FPG, HbA1-C) and/or verification of present and past clinical records of both parents. FPG and HbA1-C levels of non diabetic parents were determined to reconfirm absence of diabetes. All non diabetic parents had normal FPG (< 126 mg/dl) and HbA1-C (3.5 – 5.5%).

Three generation pedigrees were drawn to ascertain the family history and consanguinity in the parents. Consanguinity was defined as history of marriage with first cousin.

Individuals of estranged or bereaved parents, type 1 diabetic parents, were excluded from the study. Individuals with a history of Cushing' syndrome, thyrotoxicosis, or a major illness and those on medication known to affect body composition, were excluded from the study. All subjects underwent detailed medical examination. IFG was defined by fasting plasma glucose between 100–125 mg/dl (5.6–6.9 mmol/L).²⁵

Physical Measurements and Blood Collection

Body height was measured with a stadiometer to the nearest centimeter and weight to the nearest kg. Body mass index (BMI) was calculated according to the formula:

$$\text{BMI} = \text{BW (kg)} / \text{height (m)}^2$$

In all cases blood was withdrawn between 0800 – 0900 h. Two ml of venous blood was drawn from the cubital vein after overnight fasting of 12 h. Sample was added to a fluoride EDTA tube for glucose estimation.

Analytical Methods

FPG of both subjects and parents were determined. Blood glucose levels were determined by the glucose oxidase method using a commercial reagent kit (RANDOX Laboratories, Crumlin, UK) with a HumaStar 180 chemistry analyzer (Human, Wiesbaden, Germany) in duplicate. HbA1-c of the parents was estimated by affinity liquid chromatography with a D-SI Glycomat (Provalis Diagnostics, Deeside, UK).

Statistical Analysis

Numerical values were reported as mean ± standard error of means (S.E.M) and categorical variables as proportions. Chi-square test and independent sample t-test were used to compare the two groups for categorical data and numerical respectively. Age, sex and BMI adjusted logistic regression analysis was performed to determine the association between FHD (independent variable) and IFG (dependent variable), parental consanguinity (independent variable) and IFG (dependent variable). *p* value < 0.05 was considered statistically significant. All calculations were carried out with the SPSS version 16 (SPSS, Inc. Chicago, IL, USA).

Results

A total of 226 healthy non diabetic subjects of both sexes with average age (19.9 ± 0.25) years were selected. It included 151 male and 75 female subjects. There was no significant difference in the age groups of male and female subjects with and without FHD (Table 1). Parental consanguinity was found in (69%) of subjects with FHD and (29%) of the subjects without FHD (Table 2). BMI of the subjects with FHD was significantly higher ($p < 0.001$) as compared to the subjects without FHD (Table 2). BMI of the subjects with FHD and parental history of consanguinity both was higher as compared to those with no FHD and no parental history of consanguinity (Table 3). Males with a FHD had significantly higher BMI as compared to the males without FHD whereas no statistically significant difference in the BMI of the female subjects with and without FHD was found (Table 1).

Table 1: Mean \pm SEM of age, body mass index (BMI), fasting plasma glucose (FPG) and prevalence of impaired fasting glucose (IFG) and obesity in male and female subjects according to the family history of diabetes (FHD).

* $P < 0.05$ statistically significant difference between the groups with and without FHD

	FHD +ve		FHD -ve	
	Male	Female	Male	Female
N	79	45	72	30
Age	20.13 ± 0.29	18.07 ± 0.49	21.23 ± 0.43	18.83 ± 0.82
BMI	$25.69 \pm 0.82^*$	24.05 ± 1.15	21.35 ± 0.40	21.26 ± 0.34
FPG (mmol/L)	$5.34 \pm 0.07^*$	5.56 ± 0.10	4.79 ± 0.07	4.90 ± 0.12
IFG %	28 (36%)	23 (51%)	9 (12%)	6 (20%)
Obese	8 (18%)	8 (17%)	1 (2%)	0%

Similarly there was no significant difference in the BMI of the male and female subjects with FHD. The prevalence of obesity, overweight and normal weight in the group with FHD was 19%, 17% and 64% respectively while it was 2%, 7% and 9% respectively in group without FHD (Table 2). We did not find any association of BMI with FPG levels in any of the groups.

The mean FPG (5.42 ± 0.06 mmol/L) of subjects with a FHD was significantly greater ($p < 0.001$) than those without FHD (4.82 ± 0.06 mmol/L) (Table 2). Subjects with FHD and parental history of consanguinity both, had FPG levels significantly higher ($p < 0.001$) than those with no FHD and no parental history of consanguinity (Table 3). IFG was identified in 42% subjects with a FHD and in 14% without FHD (Table 2) which was higher in females (51%) as compared to males (36%) in the group with FHD (Table 1). But

Table 2: Mean ± SEM of body mass index (BMI), fasting plasma glucose (FPG) and prevalence of impaired fasting glucose (IFG), obese, overweight, non obese and parental history of consanguinity according to the family history of diabetes (FHD).

	FHD +ve n = 124	FHD-ve n = 102
Age	19.38 ± 0.31	20.56 ± 0.40
BMI	25.09 ± 0.67*	21.33 ± 0.30
Fasting Plasma Glucose (FPG) mmol/L	5.42 ± 0.06*	4.82 ± 0.06
Impaired Fasting Glucose (IFG)%	52 (42%)	15 (14.7%)
Obese	24 (19%)	2 (2%)
Over weight	21 (17%)	7 (7%)
Non obese	79 (64%)	93 (91%)
History of Consanguinity	86 (69%)	30 (29%)

*P<0.05 statistically significant difference between the groups with and without FHD

Table 3: Mean ± SEM of body mass index (BMI), fasting plasma glucose (FPG), prevalence of impaired fasting glucose (IFG) and obesity of subjects according to the family history of diabetes (FHD) & parental history of consanguinity.

	FHD +ve Parental history of Consanguinity +ve n = 86	FHD –ve Parental history of Consanguinity –ve n = 73
Age	20.05 ± 0.38	20.70 ± 0.50
BMI	25.76 ± 0.89*	21.17 ± 0.31
Fasting Plasma Glucose (FPG) mmol/L	5.61 ± 0.07*	4.87 ± 0.07
Impaired Fasting Glucose (IFG)%	43 (51%)	14 (19%)
Obese %	21 (24%)	5 (6.8%)

*P < 0.05 statistically significant

there was no difference in the prevalence of IFG between obese male and female subjects in this group. Prevalence of IFG increased to 51% in the group with FHD along with parental consanguinity as compared to 19% in those no FHD and no history of parental consanguinity (Table 3).

We found independent association of FHD with IFG in both males and females [Chi square = 13.89, (p-value = 0 < 0.001); odds ratio (OR) 13.89] by logistic regression analysis after adjusting the data for age sex and BMI. We also found significant association [Chi square = 16.31 (p-value = 0 < 0.001); odds ratio (OR) 4.21] of (FHD × parental history of consanguinity) with IFG, suggesting that there is strong interaction of FHD and parental consanguinity on IFG.

Discussion

The results of this study indicated that parental consanguinity modifies the effect of FHD on IFG. FHD and parental history of consanguinity determine the risk for impaired fasting glucose.

T₂DM is a heterogeneous and multifactorial condition due to complex interaction of genetic environmental and cultural factors. Studies identifying risk factors for T₂DM has gained momentum in the last few years due to increasing prevalence and onset of the disease at a much younger age. Genetic background of diabetes predisposes to abnormal carbohydrate and lipid metabolism.^{7,8} Studies carried out in different populations and ethnic groups reported high prevalence of obesity and metabolic abnormalities in offspring of diabetic parents, predisposing them at the risk of developing diabetes.^{11,22,21} In our study 19% of the subjects (age 15 – 25 years) with FHD are obese which is in consistent with the study reported recently from India but lower when compared with^{26,27} studies from Eastern Europe and Middle East. When parental consanguinity was also considered along with FHD the prevalence of obesity increased to 24% which showed the effect of parental consanguinity on metabolic disturbances in the offspring.

FPG levels of both male and female subjects with FHD were within normal range but significantly high-

her as compared to those without FHD. These observations are in agreement with our earlier observations in male offspring with both diabetic parents.¹³

When subjects with FHD \times parental consanguinity were compared with the subjects with no FHD \times parental consanguinity, FPG levels were significantly ($p < 0.05$) higher. This suggests disturbances of glucose metabolism are more robustly expressed in subjects with both FHD and parental consanguinity. We did not find any association between BMI and FPG levels suggesting that disturbances of glucose metabolism are independent of body fat mass. We also analyzed the data after adjusting for BMI and the results were significant indicating interaction of FHD \times parental consanguinity on fasting glucose levels.

IFG was identified in 42% of the subjects with a FHD and the prevalence of IFG was even higher (51%) when parental consanguinity and FHD were considered together.

The prevalence observed in our study is higher than that reported in Mexicans.²⁰ This suggests that cultural factors such as parental consanguinity a contributing factor for high prevalence of Prediabetes and diabetes in South East Asians. We observed a strong association of FHD \times parental consanguinity with IFG independent of obesity suggesting that individuals with FHD and parental consanguinity should be screened for Prediabetes at an early age. Similar association between FHD and IFG has been reported in Mexican children (7 – 15 years) with FHD in first degree relative.²¹ High prevalence and association of IFG with FHD and parental consanguinity suggest that defects of glucose metabolism are genetically determined.

Conclusion

In conclusion our study suggests parental consanguinity modifies the effect of FHD on IFG. FHD and parental history of consanguinity determine the risk for impaired fasting glucose in our Pakistani population. This finding could help in the screening of at risk population for T₂DM. It is therefore recommended that children with a FHD and parental consanguinity should be screened at an early age for the detection of Prediabetes. Those found with IFG should be monitored and managed with life style modifications to prevent or delay the development of T₂DM.

Future studies on a large sample size, prepubertal children and different ethnic groups need to be carried

out to further elaborate the role of FHD and parental consanguinity.

Acknowledgements

This study was supported by a research grant from the Higher Education Commission, Pakistan.

References

1. Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature*. 2001; 414: 782-87.
2. Buse JB, Polonsky KS, Burant CF. Type 2 diabetes mellitus. In: Larsen PR, Kronenberg HM, Melmed S, Polonsky KS (eds). *Williams Text Book of Endocrinology*. Philadelphia, PA: Saunders; 2003: pp 1427-67.
3. Rizvi AA. Type 2 diabetes. epidemiological trends, evolving pathogenetic concepts, and recent changes in therapeutic approach. *South Med J*. 2004; 97: 1079-87.
4. Reinehr T. Clinical presentation of type 2 diabetes mellitus in children and adolescents. *Int J Obes* 2005; 2: 105-110.
5. Elbein SC, Hoffman MD, Bragg KL, Mayorga RA. The genetics of NIDDM. *Diabetes Care*. 1994; 17: 1523-33.
6. Weijnen CF, Rich SS, Meigs JB, Krolewski AS, Warram JH. Risk of diabetes in siblings of index cases with type 2 diabetes: Implications for genetic studies. *Diabet Med*. 2002; 19: 41-50.
7. Mogren I, Lindahl B, Hogberg U. Impaired fasting glucose and impaired glucose tolerance are related to both heredity and low birth weight. *Scand J Public Health*. 2003; 31: 382-88.
8. Hauner H, Hanisch J, Bramlage P, Steinhagen-Thiessen E, Schunkert H, Jöckel KH, et al. Prevalence of undiagnosed Type-2-diabetes mellitus and impaired fasting glucose in German primary care: data from the German Metabolic and Cardiovascular Risk Project (GEMC-AS). *Exp Clin Endocrinol Diabetes*. 2008; 116: 18-25.
9. Higgins S, Gill JM, Janilionyte R, Caslake MJ, Malkova D. Physical activity, dietary intake and metabolic risk factors in non-diabetic daughter of patients with type II diabetes. *Prev Med*. 2005; 40: 145-51.
10. Ahn CW, Song YD, Nam JH, Kim DM, Woo SO, Park SW, et al. Insulin sensitivity in physically fit and unfit children of parents with type 2 diabetes. *Diabetes Med*. 2004; 21: 59-63.
11. Praveen EP, Kulshreshtha B, Khurana ML, Sahoo J, Gupta N, Kumar G, et al. Obesity and Metabolic Abnormalities in Offspring of Subjects with Diabetes Mellitus. *Diabetes Technology and Therapeutics*. 2010; 12: 723-30.
12. Praveen EP, Kulshreshtha B, Khurana ML, Sahoo J, Gupta N, Kumar G, et al. Low HDL – cholesterol

- among normal weight, normoglycemic offspring of individuals with type 2 diabetes mellitus. *Hormones (Athens)*. 2011; 10: 57-66.
13. Shahid A, Lone KP, Saeed S, Arslan M: Male offspring of both diabetic parents have higher insulin resistance and serum leptin levels compared to those with one diabetic parent. *Hormones (Athens)* 2008; 7: 313–19.
 14. Groop L, Forsblom C, Lehtovirta M, Tuomi T, Karanko S, Nissén M, et al. Metabolic consequences of a family history of NIDDM (the Botnia study): evidence for sex-specific parental effects. *Diabetes* 1996; 45: 1585-93.
 15. Anjana RM, Lakshminarayanan S, Deepa M, Farooq S, Pradeepa R, Mohan V. Parental history of type 2 diabetes mellitus, metabolic syndrome, and cardiometabolic risk factors in Asian Indian adolescents. *Metabolism*. 2009; 58: 344-50.
 16. Abu-Hasheesh MO, Abu – Samak MS, Al-Matubsi HY, Jaradeh MS, Jarrah SS, Khuzaiie RF Association of parental history of type 2 diabetes mellitus with leptin levels in Jordanian male youths. *Saudi Med J*. 2010; 30: 882-6.
 17. Guerrero-Romero F, Rodríguez-Morán M, González-Ortiz M, Martínez-Abundis E. Insulin action and secretion in healthy Hispanic – Mexican first degree relatives of subjects with type 2 diabetes. *J Endocrinol Invest*. 2001; 24: 580–86.
 18. Rodríguez-Morán M, Guerrero-Romero F. The parental phenotype of Diabetes, but not of Essential Hypertension, is linked to the development of Metabolic Syndrome in Mexican individuals. *Acta Diabetol*. 2001; 38: 87–91.
 19. Unwin N, Shaw J, Zimmet P, Alberti KG. Impaired glucose tolerance and impaired fasting glycaemia: the current status on definition and intervention. *Diabet Med*. 2002; 19: 708-23.
 20. Guerrero-Romero F, Violante R, Rodríguez-Morán M. Distribution of fasting plasma glucose and prevalence of impaired fasting glucose, impaired glucose tolerance and type 2 diabetes in the Mexican paediatric population. *Paediatr Perinat Epidemiol*. 2009; 23: 363-69.
 21. Rodríguez-Morán M, Guerrero-Romero F, Aradillas-García C, Violante R, Simental-Mendia LE, Monreal-Escalante E, Mendoza Ede L. Obesity and family history of diabetes as risk factors of impaired fasting glucose: implications for the early detection of prediabetes. *Pediatr Diabetes*. 2010; 11: 331-36.
 22. Goran MI, Bergman RN, Avila Q, Watkins M, Ball GDC, Shaibi GQ et al. Impaired glucose tolerance and reduced beta cell function in overweight Latino children with a positive family history for type 2 diabetes. *J Clin Endocrinol Metab*. 2004; 89: 207-12.
 23. Burke GL, Webber LS, Srinivasan SR, Radhakrishnamurthy B, Freedman DS, Berenson GS. Fasting plasma glucose and insulin levels and their relationship to cardiovascular risk factors in children: Bogalusa Heart Study. *Metabolism*. 1986; 35: 441–46.
 24. Ludvigsson J, Huus K, Eklöv K, Klintström R, Lahdenperä A. Fasting plasma glucose levels in healthy preschool children: effects of weight and lifestyle. *Acta Paediatr*. 2007; 96: 706–9.
 25. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Follow-up report on the diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2003; 26: 3160–67.
 26. Kelishadi R: Childhood overweight, obesity and metabolic syndrome in developing countries. *Epidemiol Rev*. 2007; 29: 62–76.
 27. Misra A, Khurana L Obesity and the metabolic syndrome in developing countries *J Clin Endocrinol Metab* 2008; 93: S9-30.