

1. Vol. 23 No. 3 (2017): AKEMU

DOI: <https://doi.org/10.21649/akemu.v23i3>

Published: 09/21/2017

Editorial

- **Granulomatous Mastitis Mimics Breast Carcinoma- A Diagnostic Concern**

Rakhshindah Bajwa

◦ [PDF](#)

Guest Editorials

- **Adolescent Reproductive Health Education**

Tayyiba Wasim

◦ [PDF](#)

Research Articles

- **Comparison of Mulligan Sustained Natural Apophyseal Glides Versus Mulligan Natural Apophyseal Glides in Mechanical Neck Pain**

Shoaib Waqas, Syed Haider Ali Shah, Usmara Zafar, Muhammad Fiaz Akhtar

◦ [PDF](#)

- **Outcome of Coverage of Post Burn Palmar Hand Contractures with Glabrous Intermediate-Thickness Plantar (ITP) Skin Graft**

Muhammad Sohail, Muhammad Hussain, Muhammad Mustehsan Bashir, Farid Ahmad Khan, Hamid Hussain Ansari, Sadia Nosheen Jan, Afzaal Bashir, Muhammad Imran

◦ [PDF](#)

- **Comparison of Kaltenborn Segmental Traction Versus Mechanical Cervical Traction for the Management of Cervical Spondylosis**

Shoaib Waqas, Muhammad Fiaz Akhtar, Ijaz Ahmad Burq, Tariq Shafi

◦ [PDF](#)

- **Preventing Child Sexual Abuse, What Parents Know?**

Marwan A Bakarman, Zobeida O Eljaaly

◦ [PDF](#)

- **Hypoglycaemic Effect of Turmeric in Alloxan-Induced Diabetic Rats**
 - [PDF](#)
- **Correlation of Urinary Neutrophil Gelatinase Associated Lipocalin with Routine Biomarkers used to Detect Kidney Injury in Systemic Lupus Erythematosus Patients with and without Nephritis**
 - [PDF](#)
- **The Study of FEV₁, FVC and FEV₁/FVC in Type 1 Diabetes Mellitus and their 11 Correlation with Duration of Type 1 Diabetes Mellitus**

Muhammad Farooq, Sheraz Saleem, Zafar Niaz, Mian Sajjad Ahmed, Qamar Rafique, Sajid Abaidullah

- [PDF](#)
- **Presence of Celiac Disease in Type 1 Diabetic Patients Presenting at Jinnah Hospital, Lahore**
 - [PDF](#)
- **Ghrelin Level in Type 2 Diabetes Mellitus and Obesity**

Mahwish Jawed, Muhammad Shahid Saeed, Adeela Shahid, Samia Jawed

- [PDF](#)
- **Association of Biochemical Derangements with Tumor Lysis Syndrome in Common Haematological Malignancies**

Asma Rasheed, Mohammad Aafaq Agha, Lubna Humayun, Mohammad Jawaid Sabzwari, Mulazim Hussain Bukhari

- [PDF](#)
- **Long Term Follow up of Multi Drug Resistance Tuberculosis Patients After 3-4 Years of Starting Second Line Anti Tubercular Drugs**
 - [PDF](#)
- **The Comparison of Therapeutic Ultrasound and Local Steroid Injection in Treatment of Mild to Moderate Carpal Tunnel Syndrome: A Randomized Controlled Trial**

Nighat Ansar, Muhammad Adeel, Sidrah Liaqat, Iqra Maqsood, Imran Ghafoor

- [PDF](#)
- **Outcome of Parotid Gland Surgery in Terms of Complications Especially the Facial nerve Palsy at Mayo Hospital, Lahore**

Muhammad Shahid Farooq, Rabia Altaf, Ahmed Uzair Qureshi, Fatima Naumeri, Abrar Ashraf Ali

- [PDF](#)
- **Effectiveness of Core Muscle Stabilization Exercises with and without Lumbar Stretching in Non-Specific Low Back Pain**

Akhlaq Ahmed, Muhammad Sharif Waqas, Muhammad Junaid Ijaz, Muhammad Adeel, Rizwan Haider,

Muhammad Izhar Ahmed

○ [PDF](#)

- **Symptomatology and Disease Conditions of Patients Visiting the Largest Tertiary Care Hospital in Punjab, Pakistan: A One-Day Point Prevalence Study**

Saira Tariq, Ali Naveed, Izza Bazigh, Zainab Pervaiz, Kanza Noor Butt, Zabreen Tahir, Rabia Ashraf, Nida Muzaffar, M Anas Tahir, Ifrah Nadeem, M Tabish Javed

○ [PDF](#)

- **Gross Morphology of Placenta in Preeclampsia**

Zahra Haider Bokhari, Mah Jabeen Muneera, S M Samee

○ [PDF](#)

- **Outcome of Segmental Tibial Defects Treated by Distraction Osteogenesis with the help of Naseer Awais (NA) External Fixator**

Syed Asif Ali, Usman Zafar Dar, Asad Ali Ch, Tayyab Shoaib, Muhammad Ali

○ [PDF](#)

- **Antibiotic Use in Infants: A Cross-Sectional Survey Assessing the Knowledge, Attitudes and Practices of Health Professionals**

Attia Anwar, Rizwan Faisal, Asad Mahmood, Huma Naz, Syed Ali Sibtain, Laiyla Shinwari

○ [PDF](#)

- **Community Based, Multicentered, Randomized, Double Blinded, Placebo controlled Clinical trial of Trachyspermum Copticum Extract Against Stable Angina**

Maaz Ahmad

○ [PDF](#)

- **Glucose Intolerance, Insulin Resistance and Obesity Risk in Children with Family History of Type II Diabetes**

Adeela Shahid, **Muhammad Shahid Saeed**, Muhammad Shoaib

○ [PDF](#)

- **Death due to Mechanical Asphyxia: - A Medico-Legal Retrospective Study at the Department of Forensic Medicine and Toxicology, KEMU, Lahore.**

Rohma Aamer, Maliha Tahir, Ayesha Mubasher, Arif Rasheed Malik

○ [PDF](#)

- **Nursing Education and Training: The Status in the Punjab Public Sector Health Services**

Ehsanullah Tarin

○ [PDF](#)

Review Article

- [Current Status of Neurology and Neuroscience Research in Pakistan](#)

Aftab Ahmad, Sadia Ashraf, Riffat Mehboob

○ [PDF](#)

Case Reports

- [Benign Cementoblastoma: A Case Report](#)

Asad Aizaz Chatha, Kirn Zulfiqar, Saman Chaudhry

○ [PDF](#)

- [A Case Report of Unusual Presence of Fetal Bone in Cervix After Induced Abortion](#)

○ [PDF](#)

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)
- [Open Journal Systems](#)
- [Make a Submission](#)
- [Browse](#)

ISSN : 2079-0694 (Online) | 2079-7192(Print) Previous ISSN : 1684-6680 (Annals of KEMC)

Developed By: IT Department, King Edward Medical University

© Platform & Workflow by: Open Journal Systems

Designed by Material Theme

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/362034312>

Glucose Intolerance, Insulin Resistance and Obesity Risk in Children with Family History of Type II Diabetes.

Article in *Annals of King Edward Medical University Lahore Pakistan* · July 2017

DOI: 10.21649/akemu.v23i3.2025

CITATION

1

READS

7

2 authors:



Adeela Shahid

Shalamar Medical and Dental College

31 PUBLICATIONS 362 CITATIONS

[SEE PROFILE](#)



Shahid Saeed

Fatima Jinnah Medical University

13 PUBLICATIONS 104 CITATIONS

[SEE PROFILE](#)

Glucose Intolerance, Insulin Resistance and Obesity Risk in Children with Family History of Type II Diabetes

Adeela Shahid,¹ Muhammad Shahid Saeed,² Muhammad Shoab³

Abstract:

Background: Genetic milieu increases the risk of diabetes; it results in disturbance of a number of metabolic pathways involved in blood glucose levels regulation and body weight. Family history of type II diabetes is one of the potential risk factors for developing diabetes.

Objective: The objective of the research was to determine early disturbance in the carbohydrate metabolism and the risk of overweight and obesity among children with family history of type II diabetes.

Materials and Methods: A total of 184 subjects (males & females) with and without family history of diabetes (FHD) were included in the study. Body weight, height and waist circumference was measured, BMI was calculated. Fasting blood glucose (FBG), 2hr oral glucose tolerance test (2hr OGTT), HbA1C and serum insulin levels were determined in duplicate, and HOMA-IR was calculated for insulin resistance index. Results: It was observed that children of type II diabetic parents had significantly higher body weight, BMI, waist circumference, FBG, HbA1C, 2hr OGTT and HOMA-IR. There was high degree of prevalence of overweight, obesity, impaired fasting glucose (IFG) and impaired 2hr oral glucose tolerance test in subjects with FHD compared to those with no FHD. There was significant association of HbA1-C, IFG, impaired 2hr-OGTT and BMI with FHD. There was no association of FBG with BMI.

Conclusion: Children of type 2 diabetic parents have metabolic disturbances at an early age resulting in glucose intolerance and insulin resistance which predisposes them at a risk of type II diabetes and obesity.

*There is no conflict of interest between authors

Introduction

Type 2 diabetes (T2D) accounts for 90% of the cases globally and is the predominant form of diabetes affecting a large number of individuals⁽¹⁾. T2D is a multifactorial and heterogeneous disease due to complicated interaction between genetics and environment to which person is exposed⁽²⁾. The epidemic of diabetes is strongly associated to the

rapidly changing life style and socioeconomic change⁽³⁾. T2DM usually has its onset after 40 years of life. In recent years there is marked decrease in age of onset of diabetes. It is a bitter reality that there is an increase in incidence of diabetes in children, adolescent and young adults in high rates. There is a high prevalence of T2D among close relations along with a high incidence among twins. It is seen that diabetes is more common in certain ethnic groups which shows a definite relation between genetic relation of occurrence of disease^(4,5).

Family history (FH) of T2D is a potential risk factor for developing diabetes. First degree non diabetic relatives are shown to have disturbance of carbohydrate and lipid metabolism⁽⁶⁾. There is

1. Professor of Physiology, Shalimar Medical & Dental College, Lahore.
2. Assistant Professor of Physiology, King Edward Medical University, Lahore.
3. Associate Professor of Physiology, Khawaja Muhammad Safder Medical College, Sialkot

Corresponding Author: Dr. Muhammad Shahid Saeed
Assistant Professor of Physiology, King Edward Medical University, Lahore.
Email: shahid.saeed1@hotmail.com

Received 13-06-2017; Accepted 10-10-2017

definitely some relationship of genes with dysfunction of beta cell and insulin resistance in these individuals⁽⁷⁾ Recent genome-wide study has further strengthened the role of complicated interaction of genetics, epigenetics, and the environmental factors in the development of T2D⁽⁸⁾. History of diabetes in pregnancy or before that is associated with poor control of blood glucose in children suggesting that exposure of fetus to disturbed intra uterine metabolic environment that may have long term effects on offspring^(9,10).

Ethnic and racial differences have also been implicated in the occurrence of T2DM. After different research it is apparent that peoples in particular area, population and ethnicity are more prone to disease. Genetic factors, intra uterine and early childhood environment are important in determining these ethnic differences. It is also reported that South Asian adults are more predisposed to T2D⁽¹¹⁾. Consumption of large amount of saturated fats have important role in decreasing insulin sensitivity, dysfunction of β -cell, increase in body weight and disturbance of glucose regulation in genetically predisposed subjects⁽¹²⁾.

Furthermore, both obesity and FH increase the risk of T2DM.⁽¹³⁾ In a certain groups, high BMI is strongly related with impaired insulin secretion, insulin resistance and disturbed glucose metabolism⁽¹⁴⁾. A study regarding the frequency of T2D in family members of diabetic subjects of black South African ethnicity, reported that 27.3% diabetic subjects had a FH⁽¹⁵⁾. The present study was carried out to determine the early disturbances of the carbohydrate metabolism and the risk of overweight and obesity among children and adolescents with family history of type II diabetes.

Materials and Methods

A selected number of 184 healthy subjects (male=84, females=100) between the 10-20 years of age were included in the study with permission from Institutional Review Board of Shalamar Institute of Health Sciences Lahore Pakistan and a written informed consent from every subject or his parents

.A study designed proforma was distributed among all participants to get all information with special focus on their family history of diabetes and any other major related disease. As far exclusion criteria is concerned subjects with history of T1DM in either of the parents, taking medications known to alter body metabolic process, with endocrine diseases like Cushing's syndrome, acromegaly, or any major disease, . Subjects with both or either of the parents not alive were also not included in the study. Parents having FBG ≥ 126 mg/dl were labeled as diabetic⁽¹⁶⁾. All subjects included in the study had fasting blood glucose levels (FBG) < 126 mg/dl and no signs of any disease.

It was a cross sectional comparative study. On the basis of detailed medical and family history of diabetes, the participants were divided into the following groups:

Healthy subjects with H/O one or both parents with type 2 diabetes (FHD) (n=124; mean age: 17.52 yrs)

Healthy subjects with no H/O type 2 diabetes in either parent (control group) (n=60; mean age: 16.68 yrs)

All the subjects underwent a detailed general physical examination, body weight, height and waist circumference were recorded in each case (17). Body height was measured by digital scale. The height was measured by wall-mounted stadiometer. Body mass index (BMI) was determined as:

$$\text{BMI} = \text{Body weight (kg)} / \text{Body height (m)}^2$$

WHO criteria for obesity and overweight was used⁽¹⁷⁾. About 4 ml of venous blood was taken after overnight fasting of 12 h. Two ml of sample was added to a fluoride EDTA tube for glucose estimation by glucose oxidase method, whereas 2 ml of sample was added to an EDTA tube for HbA1-C measurements. In order to see glucose tolerance a Glucose tolerance test (2 hr-OGTT) was done, according to the standard international procedure⁽¹⁸⁾. All subjects were advised to take food having at least 150 gm of carbohydrates for 3 days before this OGTT. The subjects were given 75 gm anhydrous

glucose dissolved in plain water and the two ml of blood samples were collected 2 h thereafter. The blood samples were used to determine glucose levels on the same day.

All tests were done by standard methods. The diabetes was reconfirmed by determining fasting blood glucose (FBG levels). Parents having FBG ≥ 126 mg/dl were labeled as diabetic. Subjects with fasting blood glucose ≥ 100 mg/dl & < 126 mg/dl were considered to have Impaired fasting glucose (IFG) and those with > 140 mg/dl blood glucose 2h after OGTT were labeled as Impaired 2 hr-OGTT⁽¹⁶⁾

Blood glucose levels were estimated by the glucose oxidase method using RANDOX Laboratories, England reagent kit. HbA1-c was done by affinity liquid chromatography method. Fasting Serum Insulin levels were determined by ELISA using commercial kits. In order to determine insulin resistance a method called HOMA-IR (Homeostasis model assessment of insulin resistance) and following equation was used.

$$\text{HOMA-IR} = \frac{\text{Insulin (Fasting)} (\mu\text{U/ml}) \times \text{glucose (Fasting)} (\text{mmol/l})}{22.5}^{(19)}$$

SPSS version 17 was used for analysis of data. All Quantitative data were expressed as mean \pm standard error of mean (SEM). Student's 't' test was used to determine significance of differences among the groups. Then it is Chi-Square test and to see the association of FHD with BMI, HbA1-C, IFG, 2hr-OGTT and HOMA-IR. p value < 0.05 was considered statistically significant.

Results

There were 124 subjects with family history of one or both parents with type 2 diabetes (FHD) with a mean age of 17.52 yrs and 60 subjects mean age 16.68 with no history of type 2 diabetes in either parent.

As far as Body weight, BMI and waist circumference of the subjects is concerned it was significantly higher in those who have family history of diabetes as compared to controls

($p < 0.05$). There was no statistically significant difference in the body height of subjects with and without family history of diabetes (Table 1).

There was higher prevalence of overweight and obese subjects (36%) with FHD compared to those with no FHD (10%) (Table 3). When the group with FHD was further analyzed 16% of the subjects were overweight (BMI > 25 & < 30) and 19% were obese (BMI > 30).

Glycosylated Hemoglobin (HbA1C), Blood Glucose Fasting (FBG) and 2hr OGTT:

Subjects with FHD had significantly higher HbA1-C, FBG and 2hr-OGTT ($P < 0.05$) compared to control group (Table 2). Large number of subjects (41%) with one or both parents with type 2 diabetes had HbA1-C $> 5.5\%$ as compare to controls (16%). There was also significant association of HbA1-C with FHD ($p < 0.0009$) (Table 3).

There was significantly higher number of subjects with IFG (29%) and impaired 2hr-OGTT (22%) in the group with FHD compared to those with no FHD. There was significant association of FHD with IFG ($p < 0.0001$) and 2hr-OGTT ($p < 0.0002$) (Table 3). We did not find any association of FBG with BMI ($p > 0.05$).

Fasting insulin levels although were high in subjects with FHD compared to control group but this difference was statistically not significant ($p > 0.05$) (Table 2)

Subjects with FHD had significantly higher HOMA-IR ($P < 0.05$) compared to control group (Table 2). Significant number of subjects (37%) with FHD had insulin resistance as shown in Table 3.

Table 1: Anthropometric Characteristics of subjects with Family History of Diabetes (FHD) and non diabetic parents (control)

Group	n	Body weight (Kg)	Height (m)	BMI Kg/m ²	Waist circumference
FHD	124	64.82±1.9 ^a	1.66±0.01	25.90±0.68 ^a	86.02±0.90 ^a
Control	60	61.48±1.5	1.68±0.01	21.49±0.42	73.11±0.86

Mean ± SEM of study variables, a significantly different from the appropriate group (P<0.05; student t-test): a FHD vs Control

Table 2: Metabolic Characteristics of subjects with FHD and non diabetic parents (control)

Group	Hb A1-c%	Glucose (mg/dl)	2hr -OGTT (mg/dl)	Fasting Insulin (µIU/L)	HOMA-IR	p value
FHD	5.63±0.05 ^a	97.76±1.16 ^a	119±2.94 ^a	12.51±1.08	3.44±0.31 ^a	0.002
Control	5.11±0.08	84.34±1.19	97.22±2.53	9.97±0.79	1.74±0.13	0.1

Mean ± SEM of study variables, a Significantly different from the appropriate group (P<0.05; student t-test): a FHD vs Control

Table 3: BMI, HbA1-C, Impaired Fasting Glucose (IFG), Impaired 2hr oral Glucose (2hr-OGTT) and HOMA-IR of the subjects with and without FHD

	FHD	Control	² X ² /P/OR
BMI>25	(36 %)	(10%)	13.9/0.0001/5.12
HbA1-C >5.5%	(41%)	(16%)	10.92/0.0009/3.4
Impaired Fasting Glucose (IFG) ≥100mg/dl & <126mg/dl	(29%)	(5%)	13.98/0.0001/7.7
Impaired 2hr oral Glucose Tolerance Test (2hr-OGTT) >140mg/dl	(22%)	(1.6%)	13.2/0.0002/17.2
HOMA-IR	(37%)	(10%)	12.5/0.0001/5.02

Percentages in parenthesis, P<0.05 statistically significant, X² = Chi-square and OR= Odds ratio

Discussion

In the present study children with history of diabetic parents had significantly high body weight, BMI and waist circumference, 36% of the subjects were found to be overweight and obese. 19% of the subjects with FHD were obese which is consistent with reports from India but lower when compared with studies from Eastern Europe and Middle East^(20,21).

FBG, HbA1C and 2hr OGTT of the children of diabetic parents was also significantly higher as compared to control group, there was high prevalence of IFG and impaired 2hr OGTT in subjects with FHD compared to those with no FHD in the current study. It was further confirmed by a contemporary study which indicated that glucose tolerance was interconnected with a family history of diabetes⁽²²⁾. Other studies have reported the fact that FHD

is positively linked to disturbance of carbohydrate metabolism associated complications and obesity in children^(5,23,24). Our data clearly demonstrates that there are dysregulated carbohydrate and lipid metabolism in these children. FHD confer a risk for metabolic disturbances such as IFG, Impaired 2hr-OGTT, insulin resistance and obesity. Parental History of T2D has been associated with vigorous expression of T2D associated risk factors in adolescents and young male adults⁽²⁵⁾. Similarly in Italian overweight and obese children high prevalence of glucose intolerance has been reported as compared to Europeans⁽²⁶⁾.

Disturbance of glucose metabolism in genetically predisposed individuals due to reduced insulin sensitivity and beta-cell function have indicated the involvement of multiple factors in patho-

physiology of Impaired Glucose Tolerance and T2D⁽²⁾. We did not find any association of FBG with BMI suggesting that disturbance of glucose metabolism is linked to genetic background and is independent of body fat mass. Whereas a recent study has reported the association of high BMI with decreased insulin sensitivity in Chinese with type 2 diabetes mellitus⁽²⁷⁾. In the current study the subjects with FHD had high HOMA-IR indicating the occurrence of insulin resistance in children. The high frequency of insulin resistance seen in adults is apparent in children at a very younger age⁽¹⁴⁾. Small sample size and failure to perform hyperinsulinemic clamp test for insulin resistance due to limited resources and facilities are the weakness of the study. However this study has provided us with markers of early disturbance of glucose regulation in healthy children of diabetic parents. Further studies on a large sample size and in children between 5 to 10 years of age should be carried out to investigate how early we must take measures to adopt healthy life style in children to prevent onset of glucose intolerance and insulin resistance.

Conclusion

The children with parental history of T2D must be monitored for glucose intolerance, insulin resistance and BMI annually. Parental history of T2D increases the risk of not only glucose intolerance and T2D but also other cardio-metabolic risk factors like overweight and obesity. It has been seen that those individuals found to have IFG and IGT, with an A1C of 5.7–6.4% suffer from insulin resistance and are more prone to diabetes as well as heart and vascular disease and they should be advised to adopt effective strategies in the form of change in life style to lower their risks.

References

1. Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature*. 2001; 414: 782-787.
2. Henninger J, Hammarstedt A, Rawshani A, Eliasson B. Metabolic predictors of impaired glucose tolerance and type 2 diabetes in a predisposed population: A prospective cohort study. *BMC Endocr Disord*. 2015; 15: 51.
3. King H, Rewers M. Diabetes in adults is now a third world problem. The WHO Ad Hoc Diabetes Reporting Group. *Bull World Health Organ*. 1991; 69: 643-648.
4. Elbein SC, Hoffman MD, Bragg KL, Mayorga RA. The genetics of NIDDM. *Diabetes Care*. 1994; 17: 1523-1533.
5. Weijnen CF, Rich SS, Meigs JB, Krolewski AS, Warram JH. Risk of diabetes in siblings of index cases with type 2 diabetes: Implications for genetic studies. *Diabet Med*. 2002; 19: 41-50.
6. Stadler M, Pacini G, Petrie J, Luger A, Anderwald C. Beta cell dysfunction in non-diabetic offspring of diabetic patients. *Diabetologia*. 2009; 52(11): 2435–2444.
7. Boesgaard TW, Gjesing AP, Grarup N, Rutanen J, Jansson PA, Hribal ML, et al. Variant near ADAMTS9 known to associate with type 2 diabetes is related to insulin resistance in offspring of type 2 diabetes patients—EUGENE2 study. *PLoS One*. 2009; 4(9): e7236.
8. Bernstein D, Golson ML, Kaestner KH. Epigenetic control of β -cell function and failure. *Diabetes Res Clin Pract*. 2017; 123: 24-36.
9. Chernauek SD, Arslanian S, Caprio S, Copeland KC, El Ghormli L, Kelsey MM, Koontz MB, Orsi CM, Wilfley D. Relationship Between Parental Diabetes and Presentation of Metabolic and Glycemic Function in Youth With Type 2 Diabetes: Baseline Findings From the TODAY Trial. *Diabetes Care*. 2016; 39(1): 110-117.
10. Miranda-Lora AL, Vilchis-Gil J, Molina-Díaz M, Flores-Huerta S, Klünder-Klünder M. Heritability, parental transmission and environment correlation of pediatric-onset type 2 diabetes mellitus and metabolic syndrome-related traits. *Diabetes Res Clin Pract*. 2017; 126: 151-159.
11. Whincup PH, Gilg JA, Owen CG, Odoki K, Alberti KG, Cook DG. British South Asians aged 13-16 years have higher fasting glucose and insulin levels than Europeans. *Diabet Med*. 2005; 22(9): 1275-1277.
12. Hu FB, van Dam RM, Liu S. Diet and risk of Type II diabetes: the role of types of fat and carbohydrate. *Diabetologia*. 2001; 44:805–817.

13. Arslanian SA. Metabolic differences between Caucasians and African-American children and the relationship to type 2 diabetes mellitus. *J Pediatr Endocrinol Metab.* 2002; 15: 509-517.
14. Whincup PH, Gilg JA, Papacosta O, Seymour C, Miller GJ, Alberti KG et al. Early evidence of ethnic differences in cardiovascular risk: Cross-sectional comparison of British South Asian and White children *BMJ.* 2002; 324 (7338): 635.
15. Erasmus RT, Blanco EB, Okesina AB, Arana JM, Gqweta Z, Matsha T. Importance of family in type 2 black South African diabetic patients. *Postgrad Med J.* 2001; 77: 323- 325.
16. American Diabetes Association. Standards of medical care in diabetes. *Diabetes Care* 2013; 36(1):S11-66.
17. World Health Organization. Measuring obesity classification and description of anthropometric data: Report on a WHO consultation of the epidemiology of obesity. Warsaw 21-23 Oct 1987 Copenhagen: WHO; 1989.
18. World Health Organization. Diabetes Mellitus: Report of a WHO study group. Geneva: World Health Organization; 1985; Tech. Rep. Ser., No. 727.
19. Matthews D.R, Hosker J.P, Rudenski A.S, Naylor B.A, Treacher D.F, Turner R.C. Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985; 28: 412-419.
20. Kelishadi R. Childhood overweight, obesity and metabolic syndrome in developing countries. *Epidemiol Rev* 2007; 29: 62–76.
21. Misra A, Khurana L. Obesity and the metabolic syndrome in developing countries. *J Clin Endocrinol Metab.* 2008; 93: S9–30.
22. Eyzaguirre F, Bancalari R, Román R, Silva R, Youlton R, Urquidi C, García H, Mericq V. Prevalence of components of the metabolic syndrome according to birthweight among overweight and obese children and adolescents. *J Pediatr Endocrinol Metab.* 2012; 25(1-2): 51-6.
23. Marcovecchio M, Mohn A, Chiarelli F. Type 2 diabetes mellitus in children and adolescents. *J Endocrinol Invest* 2005; 28:853-863.
24. Goran MI, Bergman RN, Avila Q, Watkins M, Ball GDC, Shaibi GQ et al. Impaired glucose tolerance and reduced cell function in overweight Latino children with a positive family history for type 2 diabetes. *J Clin Endocrinol Metab.* 2004; 89:207-212.
25. Shahid A., Lone KP, Saeed S, Arslan M. Male offspring of both diabetic parents have higher insulin resistance and serum leptin levels compared to those with one diabetic parent. *Hormones.* 2008; 7(4):313-319.
26. Cambuli VM, Incani M, Pilia S, Congiu T, Cavallo MG, Cossu E, Sentinelli F, Mariotti S, Loche S, Baroni MG. Oral glucose tolerance test in Italian overweight/obese children and adolescents results in a very high prevalence of impaired fasting glycaemia, but not of diabetes. *Diabetes Metab Res Rev.* 2009; 25(6): 528-534.
27. Zhao Q, Laukkanen JA, Li Q, Li G. Body mass index is associated with type 2 diabetes mellitus in Chinese elderly. *Clin Interv Aging.* 2017; 12:745-752.

Conflict of Interest : None
Funding Source: None